PROPOSED MINING FROM A PORTION OF THE REMAINING EXTENT OF THE FARM ELANDS SPRUIT 5523, LADYSMITH, KWAZULU-NATAL PROVINCE

INVASIVE PLANT SPECIES MANAGEMENT PLAN



SEPTEMBER 2022

Reference number: KZN 30/5/1/3/2/10817 MP

PREPARED FOR: Raubex Construction (Pty) Ltd P.O. Box 66192 Highveld 0169

Tel: 031 700 6411

E-mail: gary.c@raubex.com

PREPARED BY:

Greenmined Environmental (Pty) Ltd Office 107, M01 Baker Square Block De Beers Avenue Paardevlei Somerset West

7130

Tel: 021 851 2673 Fax: 086 546 0579

E -mail: christine.f@greenmined.co.za







TABLE OF CONTENTS

1.	INTRODUCTION	3
2.	OBJECTIVE	3
2	WHAT IS A PROBLEM PLANT?	2
).	WHAT IS A FROBLEW FLANT?	3
4.	WHAT TO DO WITH PROBLEM PLANTS?	4
5.	CONTROL METHODS	5
3 .	SITE SPECIFIC CONDITIONS	7
7.	PROPOSED MANAGE/CONTROL METHODS FOR MOST COMMON INVASIVE	
PL	ANT SPECIES	11





1. INTRODUCTION

The Applicant, Raubex Construction (Pty) Ltd, applied for environmental authorization to extend the existing quarry on a portion of the Remaining Extent of the farm Elands Spruit 5523 GS, Ladysmith District, KwaZulu-Natal Province.

The proposed project will entail the mining of the existing quarry pit through open-cast methods of the hard rock. The recovered material will be crushed and screened to produce gravels that can be used for road building purposes.

Although the footprint of the proposed mining area extends over a historically abandoned quarry, and therefore disturbed area, the proposed activity will still require the removal of indigenous vegetation during the site establishment- and operational phases. Due to the previous disturbance of the vegetation the presence of scattered invasive plants was identified within the footprint of the proposed mining area.

Control of invasive plant species is an important aspect during all phases of the proposed activities. Therefore, an invasive plant control plan was developed for the site to be implemented during the construction/site establishment-, operational-, decommissioning phase and 12 months' aftercare period of the mining activity.

2. OBJECTIVE

The objective of an invasive plants control plan is to provide site management with an implementation tool to control problem plant species that is present or may germinated within the proposed footprint area.

3. WHAT IS A PROBLEM PLANT?

According to the book, Problem Plants of South Africa (Bromilow 2001) a weed is a plant in the wrong place at the wrong time. Problem plants are described as vigorous growers that are easily adaptable and mostly exotic or foreign in origin. Weeds usually are pioneer plants that invade disturbed areas such as stockpile areas, overburden and topsoil stockpiles and firebreaks. Invasive plants are plants that have been imported and can invade the natural vegetation.

The National Environmental Management: Biodiversity Act, 2004 (Act No 10 of 2004) (NEMBA) came into effect in June 2004 to manage and conserve the South African biodiversity within the framework of the National Environmental Management Act, 1998



(Act No 107 of 1998) (NEMA). The Alien and Invader Species (AIS) regulations was subsequently published in terms of section 97(1) of



NEMBA in August 2014 and amended in July 2016. The AIS regulations, 2014 grouped plants into four categories and prescribes the subsequent management of each category.

- Category 1a: Invasive species which must be combatted and eradicated. Any form of trade or planting is strictly prohibited.
- Category 1b: Invasive species which must be controlled and wherever possible, removed and destroyed. Any form of trade or planting is strictly prohibited.
- * Category 2: Invasive species, or species deemed to be potentially invasive that may be grown if a permit is obtained.
- Category 3: Invasive species which may remain in prescribed areas or provinces.
 Further planting, propagation, or trade is however prohibited.

To identify invasive plants that need to be controlled/eradicated from site, the plants specified in these groups must be used as a guideline.

4. WHAT TO DO WITH PROBLEM PLANTS?

Working for Water provides the site manager with an implementation tool to control problem species and keep the site free of invasive plants:

- Step 1: Conduct Site Assessment;
- <u>Step 2</u>: Set objectives based on resources available and priorities:
 - Prioritize management of plants according to the categories stipulated in the AIS regulations.
- Step 3: Develop and implement an action plan to achieve objectives:
 - ★ The plan must be long term and should include a clearing plan that includes follow up actions for rehabilitation of the cleared area.
 - * The site plan should include a map showing the areas invested with problem plants.
 - Lighter invested areas should be cleared first to prevent the build-up of seed banks, while the control plan works progressively towards the areas with denser stands.
 - ★ Educate workers on the species that needs to be eradicated, as well as the specific method to be used.



- Conduct control of invasive plant species.
- * Remove plant remains to a suitable disposal area.
- Prevent dispersal of seeds.
- Strive for collective management and planning with neighbors to prevent seed dispersal of problem plants across boundaries.

Step 4: Monitor performance and change actions if necessary

Conduct monthly inspections to enable early detection of grow back.

5. CONTROL METHODS

The control methods to be implemented on site will depend on the specific problem plants that invaded the site. The best success is generally achieved through a combination of chemical and mechanical control methods with continuous follow-up actions. Site management must take care that the clearing methods used do not encourage further invasion through unnecessary disturbance of soil or naturally vegetated areas. The Department of Water and Sanitation's Working for Water section provides guidelines to the preferred clearing methods for most problem plants. This information can be obtained from their website: http://www.dwaf.gov.za/wfw/Control/. The selection of appropriate methods of control shall be based on the species to be controlled, the size of the plants, the density of the stand, the accessibility of terrain and environmental safety.

The Department of Water and Sanitation proposes that the following methods of control for age or size target plants:

* Seedlings

Hand pulling or hoeing:

- Hand pulling/hoeing should be carried out in sparse stands.
- Seedlings should be severed below the soil surface or removed from the soil. Soil disturbance should be minimized to reduce re-germination.

Herbicides:

Herbicides can be used on dense stands.


* Saplings

Hand pulling or hoeing:

Where appropriate saplings can be removed manually as described above.

Herbicides:





- Basal stem treatments of suitable herbicides in diesel can be carried out to the bottom 250 mm of the stem. Applications should be by means of a low pressure, coarse droplet spray from a narrow angle solid cone nozzle.
- Cut stump treatments can be used where stems are cut as low as practical. Herbicides are applied in diesel or water as recommended for the herbicide. Applications in diesel should be to the whole stump and exposed roots and in water to the cut area as recommended on the label.
- The application of herbicides should only be sprayed/used on site by a registered pest control officer.
- Mature Trees (trees above shoulder height or robust bushes 12 1 months or older)

Ring Barking:

- ➤ Bark must be removed from the bottom of the stem to a height of 0.75 1.0 m.
 All bark must be removed to below ground level for good results.
- Where clean de-barking is not possible due to crevices in the stem or where exposed roots are present, a combination of bark removal and basal stem treatment should be carried out.

Frilling or partial frilling:

Cuts should be made through the bark into the sapwood by means of a light axe and a suitable herbicide must be applied into the cuts.


Basal stem treatments:

Suitable herbicides should be applied in diesel to the base of the stem and to any exposed roots. Stems with a diameter up to 50 mm should be treated to a height of 250 mm and stems above 50 m diameter to a height of 500 mm. This method is only suitable for stems up to 100 mm in diameter.

Cut stump treatment:

X Stumps should be cut as low as practical, and the herbicide applied. Applications in diesel should be to the whole stump and exposed roots and in water to the cut area as recommended on the label.







- Herbicides selected for control shall be registered for use on that species under the conditions specified.
- * Protection of the environment is of prime importance. Riparian areas must be protected and only herbicides that are approved may be used. Washing of equipment or disposal of waste spray mixture is prohibited in or near water courses where contamination of water can occur.
- Empty herbicide containers must be disposed of as hazardous waste and may not be used for any other purpose.
- Equipment must be washed where there is no danger of contamination of a water source or natural vegetated area. It is proposed that washing be restricted to the wash bay.
- * Product and spray mixtures should be stored so that it is inaccessible to the public. Site management must ensure that the Safety Data Sheet of the product is available on site.
- * The application of herbicides should only be sprayed/used on site by a registered pest control officer.

6. SITE SPECIFIC CONDITIONS

The groundcover of the proposed mining area comprises mainly of a well-established grass layer with a moderate tree component in the previously disturbed footprint of the abandoned quarry pit.



Figure 1: Photograph of the mining area and surrounding environment looking towards the south, showing grassveld with tree component.



The previously disturbed area acts as a distribution source of invasive plant species. Lantana camara (Lantana) and Solanum mauritianum



(Bugweed) are a particular problem in the existing quarry pit, and attention should be given to the removal/control of these plants as they are both classified as Category 1b invasive plants. The Seringa trees (*Melia azedarach*), present in and around the quarry area, are also classified as Category 1b invasive plants and therefore must be controlled and where possible removed and destroyed.



<u>Figure 2</u>: View into the existing quarry pit on the property.

To prevent the dispersal of the invasive plant species to the less- or non-infected sections of the proposed mining area, site management must remove the invasive plants prior to the stripping of topsoil. A combination of the control methods, as prescribed in this plan, must be implemented during the site-establishment phase.

The entire mining footprint area, in particular the topsoil heaps, must be monitored for the duration of the operational phase as well as the first 12 months after rehabilitation of the area, to ensure the early detection and control/management of invasive plant species germinating because of vegetation removal and soil disturbance.

As everyone isn't familiar with the identification of plant species, photographs of the most important species to be controlled on site was included below for ease of reference.





Lantana

Lantana camara





Bugweed

Solanum mauritanium









<u>Syringa</u> Melia azedarach







7. PROPOSED MANAGEMENT/CONTROL METHODS FOR MOST COMMON INVASIVE PLANT SPECIES

NB: THE PROPOSED CONTROL METHODS ARE ONLY <u>RECOMMENDATIONS</u> BASED ON INFORMATION AVAILABLE TO THE ENVIRONMENTAL CONSULTANT AT THE TIME. THE ENVIRONMENTAL CONSULTANTS EMPLOYED AT GREENMINED ENVIRONMENTAL ARE NOT REGISTERED PEST CONTROL OPERATORS (PCO) AND IN THE CIRCUMSTANCES THE SITE SHOULD ENSURE THAT THE EXPERT ADVICE AND OPINION OF A REGISTERED PCO IS SOUGHT PRIOR TO THE COMMENCEMENT AND IMPLEMENTATION OF CONTROL METHODS PERTAINING TO INVASIVE SPECIES.

SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b American bramble		Foliar Spray: Confront 360 SL Plenum 160 ME Glyph 360 SL Mamba 360 SL Access 240 SL Garlon	Herbicides ★ The underground runners make the bramble very difficult to eradicate and specialised herbicides has the best results. ★ Apply as specified by supplier.





SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Black locust	5341035	Foliar Spray: Confront 360 SL Plenum 160 ME	Mechanical Control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Brazilian Pepper Tree		Foliar Spray: Lumberjack 360 SL Timbrel 360	Mechanical Control * Pull out during seedlings stage (before seed ripens) * Disposal of eradicated plants: • Dispose with general waste, • Use in areas prone to erosions, • Dispose of plant material into quarry pit. * If seeds have ripened, pull out plants while making sure seeds do not fall out. * Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. * File proof of delivery to the facility. * Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. * Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Bugweed	Photo by Phil Berdle	Foliar Spray: Confront 360 SL Starane 200 EC Tomahawk 200 Roundup Max 680 WG Tumbleweed 240 SL	Mechanical Control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 2 Castor-oil Plant		Cut Stump / Frill: Confront 360 SL Plenum 160 ME Chopper 100 SL Hatchet 100 SL	 Mechanical Control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides * Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Fountain Grass		Foliar Spray: Roundup	Herbicides ★ The species can be controlled by the usual industrial herbicides used on road sides.



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Guava		Foliar Spray: Plenum 160 ME Cut stump / Frill: Chopper 100 SL Hatchet 100 SL	Mechanical control Rull out during seedlings stage Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides.



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Inkberry		Cut Stump / Frill: Chopper 100 SL Hatchet 100 SL	 Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier





SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Lantana	(a) Exotic Plants	Foliar Spray: Plenum 160 ME Roundup Max 680 WG Tumbleweed 240 SL Glyph 360 SL Roundup Turbo 450 SL Hatchet 100 SL Mamba Max 480 SL	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier





SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Large Cocklebur	203/12/2018 11:10	Foliar Spray: Roundup	Mechanical control Rull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier





SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Large Thorn Apple	Large Thorn-Apple	Foliar Spray: Roundup These plants are annual and die once seeds have ripened. Still it is very important to remove the dead plants as leaving them will result in vigorous grow-back the following season.	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Mauritius Thorn		Foliar Spray: Roundup Max 680 WG Glyph 360 SL Mamba 360 SL Mamba Max 480 SL Roundup 360 SL Roundup Turbo 450 SL Garlon 480 EC	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Mexican Poppy	Mexican Poppy	Foliar Spray: Access 240 SL These plants are annual and die once seeds have ripened. Still it is very important to remove the dead plants as leaving them will result in vigorous grow-back the following season.	 Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Mexican Sunflower		Foliar Spray: Grazon DS {triclopyr (300 g/L) + picloram (100 g/L)}	Mechanical control Dull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Paraffin / Triffid weed		Foliar Spray: Confront 360 SL Plenum 160 ME Roundup Max 680WG Glyph 360 SL	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Peanut Butter Cassia		Cut Stump / Frill: Chopper 100 SL Hatchet 100 SL	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier





SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Queen of the Night		Full Cover Spray: MSMA 720 SL	Mechanical control Plants can be chopped down, but the stem base must be dug up. Cut plants must be deeply buried or left on a cemented area until dried out. Chopped or broken branches are capable of taking root and forming new plants, therefore pieces should not be carted away or discarded. Herbicides Small plants can be sprayed and larger ones injected with MSMA.





SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Red Sesbania	Sesbania punicea	Foliar Spray (Seedlings <1m): Roundup 360 SL Roundup Max 680 WG Glyph 360 SL Mamba 360 SL Mamba Max 480 SL Springbok 360 Foliar Spray (Seedlings 1 - 2 m): Roundup Max 680 WG Glyph 360 SL Mamba 360 SL Roundup 360 SL Garlon 4 EC Foliar Spray (Adult): Roundup Max 680 WG Chopper 100 SL Hatchet 100 SL	Mechanical control * Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier





SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Spear Thistle / Scotch Thistle	Shear Thisle	Foliar Spray: Confront 360 SL Plenum 160 ME Access 240 SL These plants are annual and die once seeds have ripened. Still it is very important to remove the dead plants as leaving them will result in vigorous grow-back the following season.	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier





SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Seringa		Cut stump / Frill: Confront 360 SL Plenum 160 ME Chopper 100 SL Hatchet 100 SL Access 240 SL	Mechanical control Pull out during seedlings stage Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Thorned Bitter Apple	T-porned Elver Apple	Foliar Spray: Roundup Access 240 SL	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1a – 3 Wattle species	Black Wattle	Seedlings – 1m Foliar Spray: Confront 360 LS Starane 200 EC Tomahawk 200 EC Volvoxypyr 200 EC Roundup 360 SL RoundUp Turbo 450 SL	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Wild Tobacco	Eintronmentall Wood Wild Tobadeb	Foliar Spray: Access 240 SL 2,4,5- trichlorophenoxyacetic acid (2,4,5-T)	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



SPECIES	РНОТО	HERBICIDES	CONTROL METHOD
Category 1b Yellow Firethorn	Distribution of the Literature of the Control of th	Cut Stump / Frill: Access	Mechanical control Pull out during seedlings stage (before seed ripens) Disposal of eradicated plants: Dispose with general waste, Use in areas prone to erosions, Dispose of plant material into quarry pit. If seeds have ripened, pull out plants while making sure seeds do not fall out. Place plants in a black plastic bag and dispose of at an incineration facility to be destroyed. File proof of delivery to the facility. Alternatively, the removed plants can be buried in a trench of at least 1m deep. Grow-back will need to be controlled in this area, preferably with herbicides. Herbicides Apply as specified by supplier



8. REFERENCES

Bromilow C. 2001. Problem Plants of South Africa. Briza Publications. South Africa

Department of Environmental Affairs. 2016. Alien and Invasive Species List, 2016. Cape Town: Government Printer

Invader Plant Specialists (Pty) Ltd. 2017. Herbicide Applicators Course. Durbanville

South Africa. 2004. National Environmental Management: Biodiversity Act No 10 of 2004. Cape Town: Government Printer.

Todd S. 2012. Alien Invasive Plant Management Plan: Solar Direct Graspan Solar Energy Facility

Working for Water. Indigenous Replacement Plants in the KZN region. Department of Water Affairs. South Africa

Working for Water. 2002. Policy on the Use of Herbicides for the Control of Alien Vegetation. Department of Water Affairs. South Africa

